

**Q20: From what you have seen or heard...which of the following results in the highest number of deaths in Ohio each year?**

		TRAFFIC ACCIDENTS	FALLS	FIREARMS	ACCIDENTAL DRUG POISONINGS AND DRUG OVERDOSES	DON'T KNOW / NOTSURE	Count
Ohio Residents		48.6%	3.6%	13.4%	31.5%	2.8%	945
Sex	Male	49.5%	2.9%	12.1%	33.1%	2.5%	455
	Female	47.9%	4.2%	14.6%	30.1%	3.2%	490
Sex and Race Combined Variable	African-American	40.8%	.6%	39.2%	18.5%	.8%	100
	White male	51.3%	3.6%	9.4%	33.2%	2.5%	367
	White female	50.7%	4.9%	9.6%	31.9%	2.9%	408
Age of Respondent	18 to 29	47.4%	1.9%	16.1%	33.1%	1.4%	187
	30 to 45	48.2%	2.7%	14.8%	32.7%	1.5%	261
	46 to 64	50.0%	5.6%	10.1%	31.8%	2.5%	312
	65 and over	47.7%	3.4%	14.6%	27.2%	7.1%	170
Education	Less than high school	36.3%	1.5%	22.2%	38.8%	1.3%	119
	High school graduate	40.9%	2.6%	15.5%	35.6%	5.4%	330
	Some college	51.5%	4.8%	12.1%	31.0%	.6%	282
	College graduate	64.1%	4.8%	7.0%	21.4%	2.7%	211
Poverty Status	100% or below FPL	30.3%	.8%	20.7%	45.5%	2.6%	142
	Between 100% and 200% FPL	52.8%	2.8%	11.6%	31.6%	1.2%	142
	Above 200% FPL	52.2%	3.8%	10.8%	31.0%	2.1%	423
Registered to Vote	Yes	49.2%	4.3%	13.9%	30.8%	2.0%	759
	No	46.8%	.8%	11.7%	34.2%	6.4%	185
Party ID	Democrat	45.5%	3.8%	19.4%	28.6%	2.7%	367
	Independent	52.2%	5.7%	7.5%	32.1%	2.5%	152
	Republican	51.4%	2.9%	10.6%	32.2%	2.9%	334
Insurance Status	Insured	49.9%	3.9%	12.3%	30.8%	3.2%	805
	Not Insured/Don't Know	42.5%	1.9%	18.7%	36.3%	.6%	136
Area of Dominant TV Influence	Cincinnati	47.3%	1.5%	16.8%	31.0%	3.3%	138
	Cleveland	55.2%	3.1%	10.2%	28.3%	3.3%	306
	Columbus	42.5%	4.1%	17.8%	32.0%	3.5%	190
	Dayton	53.6%	7.4%	6.4%	30.1%	2.5%	115
	Other	42.7%	3.1%	15.8%	37.0%	1.4%	194

**Q20: From what you have seen or heard...which of the following results in the highest number of deaths in Ohio each year?**

		<b>TRAFFIC ACCIDENTS</b>	<b>FALLS</b>	<b>FIREARMS</b>	<b>ACCIDENTAL DRUG POISONINGS AND DRUG OVERDOSES</b>	<b>DON'T KNOW / NOTSURE</b>	<b>Count</b>
Geographical Region	Northeast Ohio	51.2%	3.1%	11.5%	31.3%	2.9%	375
	Northwest Ohio	48.7%	5.1%	15.6%	29.9%	.7%	118
	Central Ohio	44.2%	5.1%	21.0%	25.4%	4.3%	155
	Southeast Ohio	37.9%	.2%	4.3%	56.3%	1.3%	63
	Southwest Ohio	50.7%	3.7%	12.8%	29.6%	3.2%	231
Rural or Urban County	Urban county	48.1%	3.8%	21.3%	24.0%	2.9%	439
	Suburban county	50.2%	5.7%	6.8%	33.5%	3.8%	240
	Rural, small city	48.4%	1.3%	6.3%	42.0%	1.9%	263
General Health Rating	Excellent/Very Good	48.9%	3.8%	11.2%	32.9%	3.1%	491
	Good	47.4%	4.6%	13.1%	32.4%	2.6%	292
	Fair/Poor	50.9%	1.0%	20.7%	25.0%	2.3%	158
Type of Insurance	Employer based	56.2%	4.2%	9.0%	28.1%	2.5%	484
	Medicare	43.4%	3.8%	10.4%	35.8%	6.6%	129
	Other	36.5%	3.2%	22.4%	35.0%	2.9%	187
Uninsured, Medicare, or other source of health insurance	Uninsured	42.5%	1.9%	18.7%	36.3%	.6%	136
	Medicare alone or combo	43.6%	3.4%	12.0%	35.2%	5.9%	171
	Other Source	51.1%	4.1%	12.5%	29.8%	2.5%	629
Uninsured, Medicaid, or other source of Health Insurance	Uninsured	42.5%	1.9%	18.7%	36.3%	.6%	136
	Medicaid alone or combo	28.4%		30.3%	37.7%	3.6%	94
	Other Source	52.3%	4.4%	10.0%	30.1%	3.2%	705
Abused Non-Prescribed Pain Reliever	Yes	47.6%		23.7%	28.6%		56
	No	48.8%	3.8%	12.8%	31.5%	3.0%	884
Family Member Abused Pain Relievers	Yes	38.3%	1.9%	17.7%	41.4%	.7%	192
	No	50.8%	4.1%	12.2%	29.5%	3.4%	740

Ohio Poll #125, July 2011  
 Institute for Policy Research  
 University of Cincinnati  
 The Health Foundation of Greater Cincinnati

**Q16: During the past five years...has a doctor, dentist, or other medical professional prescribed a pain reliever for you that you could not purchase 'over the counter'....such as oxycontin, vicodin, percocet, or codeine?**

		YES	NO	DON'T KNOW / NOT SURE	Count
Ohio Residents		54.7%	45.1%	.2%	943
Sex	Male	50.8%	49.0%	.2%	454
	Female	58.4%	41.5%	.2%	489
Sex and Race Combined Variable	African-American	43.2%	56.8%		100
	White male	54.0%	45.7%	.3%	366
	White female	59.6%	40.2%	.2%	407
Age of Respondent	18 to 29	58.5%	41.5%		188
	30 to 45	56.4%	43.6%		261
	46 to 64	56.5%	43.1%	.4%	310
	65 and over	45.7%	54.0%	.4%	171
Education	Less than high school	62.8%	37.2%		119
	High school graduate	50.5%	49.2%	.3%	331
	Some college	57.3%	42.5%	.2%	281
	College graduate	53.0%	46.9%	.1%	211
Poverty Status	100% or below FPL	64.0%	36.0%		142
	Between 100% and 200% FPL	49.8%	49.8%	.4%	142
	Above 200% FPL	55.2%	44.8%		422
Registered to Vote	Yes	57.5%	42.4%	.1%	757
	No	43.3%	56.2%	.5%	185
Party ID	Democrat	50.8%	49.2%		366
	Independent	59.9%	39.1%	1.0%	152
	Republican	56.0%	43.9%	.1%	333
Insurance Status	Insured	54.9%	45.0%	.1%	804
	Not Insured/Don't Know	54.5%	44.8%	.7%	136
Area of Dominant TV Influence	Cincinnati	58.1%	41.9%		136
	Cleveland	57.0%	43.0%		305
	Columbus	54.5%	45.3%	.1%	189
	Dayton	54.7%	44.8%	.5%	115
	Other	49.8%	49.7%	.5%	195

**Q16: During the past five years...has a doctor, dentist, or other medical professional prescribed a pain reliever for you that you could not purchase 'over the counter'....such as oxycontin, vicodin, percocet, or codeine?**

		YES	NO	DON'T KNOW / NOT SURE	Count
Geographical Region	Northeast Ohio	54.8%	45.2%		375
	Northwest Ohio	53.6%	46.4%		118
	Central Ohio	52.0%	47.8%	.2%	155
	Southeast Ohio	52.7%	45.9%	1.5%	63
	Southwest Ohio	58.3%	41.4%	.3%	230
Rural or Urban County	Urban county	52.4%	47.4%	.2%	438
	Suburban county	59.2%	40.8%		240
	Rural, small city	55.2%	44.5%	.4%	263
General Health Rating	Excellent/Very Good	49.1%	50.6%	.2%	490
	Good	54.0%	45.8%	.2%	293
	Fair/Poor	73.9%	26.1%		159
Type of Insurance	Employer based	56.7%	43.2%	.1%	482
	Medicare	52.1%	47.4%	.5%	130
	Other	50.8%	49.2%		187
Uninsured, Medicare, or other source of health insurance	Uninsured	54.5%	44.8%	.7%	136
	Medicare alone or combo	50.3%	49.3%	.4%	171
	Other Source	55.8%	44.2%	.0%	627
Uninsured, Medicaid, or other source of Health Insurance	Uninsured	54.5%	44.8%	.7%	136
	Medicaid alone or combo	49.8%	50.2%		94
	Other Source	55.2%	44.6%	.1%	704
Obesity Scale Distributions	Not Obese	55.2%	44.8%		340
	Mildly Obese	50.6%	48.9%	.5%	330
	Moderately Obese	57.8%	42.0%	.2%	154
	Severely Obese	55.6%	44.4%		45
	Very Severely Obese	74.1%	25.9%		35

Ohio Poll #125, July 2011  
 Institute for Policy Research  
 University of Cincinnati  
 The Health Foundation of Greater Cincinnati

**Q18: Have you ever, even once, used a pain reliever such as oxycontin, vicodin, percocet, or codeine...when they were not prescribed for you or that you took only for the experience or feeling they caused?**

		YES	NO	DON'T KNOW	Count
Ohio Residents		5.9%	93.5%	.5%	946
Sex	Male	7.8%	91.2%	1.0%	455
	Female	4.2%	95.7%	.1%	490
Sex and Race Combined Variable	African-American	5.1%	94.9%		100
	White male	8.6%	91.0%	.4%	367
	White female	3.6%	96.3%	.1%	409
Age of Respondent	18 to 29	16.2%	83.8%		188
	30 to 45	5.4%	94.6%		261
	46 to 64	3.4%	96.6%		312
	65 and over	.5%	96.7%	2.8%	171
Education	Less than high school	10.7%	85.6%	3.7%	119
	High school graduate	2.9%	97.0%	.1%	331
	Some college	8.7%	91.3%		282
	College graduate	4.4%	95.6%		211
Poverty Status	100% or below FPL	9.1%	89.8%	1.1%	142
	Between 100% and 200% FPL	8.3%	91.7%		142
	Above 200% FPL	4.4%	95.6%		422
Registered to Vote	Yes	5.5%	93.9%	.6%	759
	No	7.9%	92.1%		185
Party ID	Democrat	4.2%	95.8%		366
	Independent	9.9%	90.1%		152
	Republican	5.9%	92.6%	1.5%	335
Insurance Status	Insured	4.9%	94.5%	.6%	806
	Not Insured/Don't Know	12.5%	87.5%		136
Area of Dominant TV Influence	Cincinnati	7.3%	91.6%	1.1%	138
	Cleveland	4.0%	95.0%	.9%	305
	Columbus	9.4%	90.6%		190
	Dayton	7.2%	92.4%	.4%	115
	Other	4.0%	96.0%		195

**Q18: Have you ever, even once, used a pain reliever such as oxycontin, vicodin, percocet, or codeine...when they were not prescribed for you or that you took only for the experience or feeling they caused?**

		<b>YES</b>	<b>NO</b>	<b>DON'T KNOW</b>	<b>Count</b>
Geographical Region	Northeast Ohio	4.1%	95.2%	.8%	375
	Northwest Ohio	7.7%	92.3%		118
	Central Ohio	8.2%	91.8%		155
	Southeast Ohio	1.4%	98.6%		64
	Southwest Ohio	7.9%	91.2%	.9%	231
Rural or Urban County	Urban county	7.7%	91.9%	.4%	439
	Suburban county	5.6%	94.4%		240
	Rural, small city	3.3%	95.5%	1.3%	264
General Health Rating	Excellent/Very Good	3.3%	96.7%		491
	Good	9.2%	89.8%	1.0%	293
	Fair/Poor	8.1%	90.7%	1.3%	159
Type of Insurance	Employer based	5.4%	94.6%		483
	Medicare	4.4%	94.0%	1.6%	130
	Other	4.0%	94.5%	1.5%	187
Uninsured, Medicare, or other source of health insurance	Uninsured	12.5%	87.5%		136
	Medicare alone or combo	3.3%	95.5%	1.2%	171
	Other Source	5.3%	94.2%	.5%	629
Uninsured, Medicaid, or other source of Health Insurance	Uninsured	12.5%	87.5%		136
	Medicaid alone or combo	7.7%	89.3%	3.0%	94
	Other Source	4.5%	95.2%	.3%	706
Obesity Scale Distributions	Not Obese	10.6%	89.4%		341
	Mildly Obese	1.2%	98.3%	.5%	330
	Moderately Obese	9.6%	88.6%	1.8%	154
	Severely Obese		100.0%		45
	Very Severely Obese	2.2%	97.8%		35

Ohio Poll #125, July 2011  
 Institute for Policy Research  
 University of Cincinnati  
 The Health Foundation of Greater Cincinnati

**Q19: Thinking about your friends and family...have any of your family members or friends experienced problems as a result of abusing prescription pain relievers such as oxycontin, vicodin, percocet, or codeine?**

		YES	NO	DON'T KNOW	Count
Ohio Residents		20.3%	78.4%	1.3%	945
Sex	Male	20.7%	77.8%	1.5%	455
	Female	20.0%	78.9%	1.1%	490
Sex and Race Combined Variable	African-American	12.1%	86.2%	1.7%	100
	White male	19.8%	78.3%	1.9%	367
	White female	21.6%	77.4%	1.0%	408
Age of Respondent	18 to 29	29.5%	68.1%	2.4%	188
	30 to 45	19.3%	79.5%	1.2%	261
	46 to 64	22.6%	76.5%	.9%	312
	65 and over	9.0%	90.0%	1.0%	170
Education	Less than high school	29.6%	70.4%		119
	High school graduate	19.7%	78.8%	1.5%	330
	Some college	23.6%	74.4%	2.0%	282
	College graduate	11.8%	87.3%	.9%	211
Poverty Status	100% or below FPL	24.7%	74.2%	1.0%	142
	Between 100% and 200% FPL	32.8%	66.9%	.3%	142
	Above 200% FPL	17.7%	81.4%	.9%	422
Registered to Vote	Yes	20.3%	78.1%	1.6%	759
	No	20.4%	79.4%	.2%	185
Party ID	Democrat	22.3%	75.7%	2.0%	366
	Independent	23.7%	74.9%	1.4%	152
	Republican	20.1%	79.6%	.4%	334
Insurance Status	Insured	17.7%	81.7%	.6%	805
	Not Insured/Don't Know	36.6%	57.9%	5.6%	136
Area of Dominant TV Influence	Cincinnati	19.4%	78.8%	1.8%	138
	Cleveland	17.0%	82.5%	.5%	305
	Columbus	21.0%	78.1%	.8%	190
	Dayton	19.0%	79.3%	1.7%	115
	Other	26.7%	70.8%	2.5%	194

**Q19: Thinking about your friends and family...have any of your family members or friends experienced problems as a result of abusing prescription pain relievers such as oxycontin, vicodin, percocet, or codeine?**

		YES	NO	DON'T KNOW	Count
Geographical Region	Northeast Ohio	17.7%	81.8%	.5%	375
	Northwest Ohio	29.8%	70.2%		117
	Central Ohio	20.7%	78.5%	.8%	155
	Southeast Ohio	22.4%	70.1%	7.6%	64
	Southwest Ohio	19.2%	78.9%	1.9%	231
Rural or Urban County	Urban county	19.2%	79.5%	1.2%	439
	Suburban county	16.1%	83.0%	.9%	240
	Rural, small city	26.2%	72.0%	1.8%	263
General Health Rating	Excellent/Very Good	17.6%	81.7%	.7%	491
	Good	19.7%	77.7%	2.6%	292
	Fair/Poor	29.7%	69.2%	1.1%	159
Type of Insurance	Employer based	19.8%	79.7%	.5%	483
	Medicare	11.6%	87.3%	1.0%	130
	Other	17.1%	82.3%	.6%	187
Uninsured, Medicare, or other source of health insurance	Uninsured	36.6%	57.9%	5.6%	136
	Medicare alone or combo	9.2%	89.8%	1.0%	171
	Other Source	20.2%	79.3%	.5%	628
Uninsured, Medicaid, or other source of Health Insurance	Uninsured	36.6%	57.9%	5.6%	136
	Medicaid alone or combo	25.7%	74.3%		94
	Other Source	16.8%	82.5%	.7%	706
Abused Non-Prescribed Pain Reliever	Yes	53.2%	46.8%		56
	No	18.0%	80.6%	1.4%	884

Ohio Poll #125, July 2011  
 Institute for Policy Research  
 University of Cincinnati  
 The Health Foundation of Greater Cincinnati